DRUG CHEMISTRY SECTION TECHNICAL PROCEDURE MANUAL			
Procedure D-08	Extraction and Separations	_	
	Extraction of Psilocybe Mushrooms		
Effective Date:	November 20, 2006	Page 1 of 2	

Name of Procedure:

Extractions and Separations
Extraction of Psilocybe Mushrooms

Suggested Uses:

This procedure is used to extract psilocybe mushrooms.

Apparatus Needed to Perform Procedure Including Preparation of Reagent:

Fume hood Eye protection Gloves Laboratory coat Small beaker Pipets, glass, disposable Pipet bulb Glass stirring rod pH Test paper Test tube Test tube rack Clean glass vial with cap Glacial Acetic Acid Ammonium Hydroxide Ethyl Ether Nitrogen source

Formula for Preparing Reagent:

3:1 Chloroform/Isopropanol Reagent

- 1. Measure out 60 milliliters of chloroform.
- 2. Measure out 20 milliliters of isopropanol.
- 3. Combine the chloroform and isopropanol.
- 4. Pour solution into a reagent bottle.
- 5. Properly label reagent bottle.

DRUG CHEMISTRY SECTION TECHNICAL PROCEDURE MANUAL			
Procedure D-08	Extraction and Separations		
Extraction of Psilocybe Mushrooms			
Effective Date:	November 20, 2006	Page 2 of 2	

Application of Procedure on Evidence:

- 1. Break up approximately 1-2 grams of psilocybe mushrooms and place in a small beaker.
- 2. Add enough deionized water to moisten.
- 3. Add 1-2 milliliters of glacial acetic acid and check with test paper to assure that the solution is acidic.
- 4. Stir 1-2 minutes. [Do not leave in acidic solution for extended period of time]
- 5. Decant liquid to test tube.
- 6. Add concentrated ammonium hydroxide dropwise until a pH of 8 is obtained.
- 7. Gently extract with ethyl ether or 3:1 chloroform/isopropanol.
- 8. Solvent may be dried using magnesium sulfate of sodium sulfate.
- 9. Evaporate solvent under dry nitrogen.

Safety Concerns:

Ethyl ether is extremely flammable. Ammonium hydroxide is a strong base and glacial acetic acid is a strong acid. Care should be taken to keep these two components capped when not in use and away from each other to avoid mixing.

Literature References:

Casale, J.,

An Aqueous-Organic Extraction Method for the Isolation and Identification of Psilocin from Hallucinogenic Mushrooms

,

Journal of Forensic Science, January 1985.

Modified by Chemist T.H. McSwain with the North Carolina State Bureau of Investigation Drug Chemistry Laboratory, in use in the laboratory since January, 1985.