

## **Technical Procedure for Ninhydrin-HFE7100**

**1.0 Purpose** - This procedure describes how to make the Ninhydrin-HFE7100 solution and apply it to items of evidence.

**2.0 Scope** - This procedure applies to porous items of evidence that are to be examined for the presence of latent prints. Ninhydrin reacts to the amino acids present in fingerprint residue and turns the amino acids purple when the reaction is complete.

**3.0 Definitions** – N/A

### **4.0 Equipment, Materials and Reagents**

#### **4.1 Equipment and Materials**

- Laboratory coat and gloves
- Face shield visor and/or safety goggles
- Magnetic stirrer, magnetic follower, and magnetic retriever
- Glass beakers
- Graduated cylinders
- Dark, shatter-proof container
- Forceps
- Fume hood
- Glass tray, paint brush, or aerosol sprayer (for application)
- Camera/scanner
- Dust or mist respirator (for application outside of fume hood)

#### **4.2 Reagents**

- Ninhydrin crystals (25 g)
- Ethanol (ethyl alcohol) (225 mL)
- Ethyl acetate (10 mL)
- Glacial acetic acid (25 mL)
- HFE-7100 (1000 mL)

### **5.0 Procedure**

#### **5.1 Chemical Preparation**

##### **5.1.1 Concentrate Solution**

**5.1.1.1** Place twenty-five (25) grams of ninhydrin crystals and a magnetic follower into a five hundred (500) mL beaker.

**5.1.1.2** Add two hundred twenty-five (225) mL of ethanol to the beaker and stir until the ninhydrin crystals have completely dissolved. Do not use heat.

**5.1.1.3** Add ten (10) mL of ethyl acetate while stirring.

5.1.1.4 Add twenty-five (25) mL of glacial acetic acid to the mixture. Continue stirring until clear yellow solution is produced.

5.1.1.5 Transfer the concentrate solution to a clean, dark, shatterproof container.

## 5.1.2 Working Solution

5.1.2.1 Measure fifty-two (52) mL of the concentrate solution and place in a two (2) L glass beaker with a magnetic follower.

5.1.2.2 Measure one thousand (1000) mL of HFE-7100 and add to the concentrate solution while stirring.

5.1.2.3 Transfer the resulting working solution to a clean, dark, shatterproof container.

## 5.2 Processing Procedures

### 5.2.1 Chemical Application

5.2.1.1 Forensic Scientists shall produce a self-made test print to be processed concurrently with items of evidence (see Section Technical Procedure for Ensuring Quality Control).

5.2.1.2 **Dipping Method** – Completely submerge the item in the working solution for five (5) to ten (10) seconds.

5.2.1.3 **Brush Method** – Dip the brush into the working solution and brush directly onto the item.

5.2.1.4 **Spray Method** – Spray the item with the working solution to completely saturate the item.

5.2.1.5 Allow the item to dry completely prior to proceeding.

5.2.2 Latent impressions will develop over time at room temperature. Several methods are available to enhance the development process.

5.2.2.1 **Plastic Bag** – Place the item in a sealed plastic bag until latent impressions develop.

5.2.2.2 **Steam Iron** – The iron is used to provide heat and moisture to the item. Hold the iron above the item and steam it, taking care to avoid contact between the item and the iron.

5.2.2.3 **Microwave** – Heat a tray or beaker of water in the microwave to produce steam. Place the item in the microwave for approximately five (5) minutes, or until impressions develop. Do not turn on the microwave with the evidence inside and do not allow the evidence to contact the hot water bath.

5.2.2.4 **Humidity Chamber** – Large items may be placed in a humidity chamber for approximately four (4) to five (5) hours. The chamber shall be checked periodically to ensure adequate moisture is present.

- 5.2.3 Preservation of Developed Impressions** – Preserve the developed impressions through photography (see photographic equipment procedures) and/or by electronic recording (see Section Image Processing Procedure).

**5.3 Standards and Controls** – N/A

**5.4 Calibration** – N/A

**5.5 Sampling** – N/A

**5.6 Calculations** – N/A

**5.7 Uncertainty of Measurement** – N/A

## **6.0 Limitations**

- 6.1** The ninhydrin crystals shall be stored in the original shipping container until needed.
- 6.2** Ninhydrin solutions shall be stored in dark, shatter-proof containers at all times to avoid direct exposure to sunlight.
- 6.3 Shelf Life**
- 6.3.1** Ninhydrin crystals – indefinite.
- 6.3.2** Concentrate and working solution - one (1) year.

## **7.0 Safety**

- 7.1** The process shall be performed in a fume hood as the fumes may cause some irritation when in contact with the eyes or skin and may be harmful if inhaled or ingested.
- 7.2** Protective goggles, gloves and lab coats shall be worn during processing as the solution will stain skin and clothing.

## **8.0 References**

Almog, J. "Reagents for Chemical Development of Latent Fingerprints: Vicinal Triketones – Their Reaction with Amino Acids and with Latent Fingerprints on Paper." *Journal of Forensic Sciences*. Vol. 32, 6: 1565–1573 (1987).

Almog, J., A. Hirshfeld and J.T. Klug. "Reagents for the Chemical Development of Latent Fingerprints: Synthesis and Properties of Some Ninhydrin Analogues." *Journal of Forensic Sciences*. Vol. 27, 4: 912–917 (1982).

Beaudoin, A. "Research on Transferring a Fingerprint to a Ninhydrin-Treated Document." *Journal of Forensic Identification*. Vol. 54, 2: 178–184 (2004).

Burns, Gerry F.F.S. "Ninhydrin Longevity." *Fingerprint Whorld*. (April 1982): 98-100.

Hamm, E. "Enhancement and Development of Blood Prints." *74<sup>th</sup> Annual Educational Conference International Association for Identification*. (June 22, 1989): 1-11.

Herod, D.W., and E.R. Menzel. "Laser Detection of Latent Fingerprints: Ninhydrin." *Journal of Forensic Sciences*. Vol. 27, 1: 200–204 (1982).

Herod, D.W., and E.R. Menzel. "Laser Detection of Latent Fingerprints: Ninhydrin Followed by Zinc Chloride." *Journal of Forensic Sciences*. Vol. 27, 3: 513–518 (July 1982).

Kent, T., ed. *Manual of Fingerprint Development Techniques: A Guide to the Selection and Use of Processing for the Development of Latent Fingerprints*. Police Scientific Development Branch, London (July 1992).

Kent, T., D. Hewlet and V. Sears. A "Green" Formulation for Ninhydrin: Search for a Safe Effective Replacement for CFC's. Fingerprint Research Group, Police Scientific Development Branch.

Kobus, H.J., M. Stoilovic and R.N. Warrenner. "A Simple Luminescent Post-Ninhydrin Treatment for The Improved Visualization of Fingerprints on Documents in Cases Where Ninhydrin Alone Gives Poor Results." *Forensic Science International*. Vol. 22: 161–170 (1983).

Lee, H.C. "Methods of Latent Print Development." *Proceedings of the International Forensic Symposium on Latent Prints*. (July 1987): 15–24.

Lennard, C.J. and P.A. Margot. "Sequencing of Reagents for the Improved Visualization of Latent Fingerprints." *Proceedings of the International Forensic Symposium on Latent Prints*. (July 1987): 141-142.

Lennard, C.J., et al. "Synthesis and Evaluation of Ninhydrin Analogues as Reagents for the Development of Latent Fingerprints on Paper Surfaces." Forensic Science Research Unit, Chemistry Department, Australian National University.

*Manual of Fingerprint Development Techniques: A Guide to the Selection and Use of Processes for the Development of Latent Fingerprints*. Scientific Research and Development Branch, London (1986).

Menzel, E.R., R.A. Bartsch and J.L. Hallman. "Fluorescent Metal-Rhuemann's Purple Coordination Compounds: Applications to Latent Fingerprint Detection." *Journal of Forensic Sciences*. Vol. 35, 1: 25–34 (1990).

Robertson, J. *Report on Comparison of DFO with Ninhydrin*. Australian Federal Police. (1990): 3.

Schiltz, E., K.D. Schnackerz and R.W. Gracy. "Comparison of Ninhydrin, Fluorescamine, and O-Phthaldehyde for the Detection of Amino Acids and Peptides and Their Effects on the Recovery and Composition of Peptides from Thin-Layer Fingerprints." *Analytical Biochemistry*. Vol. 79: 33–41 (1977).

Stoilovic, M., and R.N. Warrenner. "Cadium Complexation: An Improved Fluorescent Technique for the Enhancement of Weak Ninhydrin-Developed Latent Fingerprints." *International Forensic Symposium on Latent Prints*. (July 1987): 157-160.

Stoilovic, M., R.N. Warrenner and H.J. Kobus. "An Evaluation of the Reagent NBD Chloride for the Production of Luminescent Fingerprints on Paper: II. A Comparison with Ninhydrin." *Forensic Science International*. Vol. 24: 279–284 (1984).

Trozzi, T.A., R.L. Schwartz and M.L. Hollars. *Processing Guide for Developing Latent Prints*. (2000): 1-64.

US Department of Justice. *Chemical Formulas and Processing Guide for Developing Latent Prints*. FBI Laboratory Division, Latent Fingerprint Section (1994).

**9.0 Records – N/A**

**10.0 Attachments – N/A**

| Revision History |                |                   |
|------------------|----------------|-------------------|
| Effective Date   | Version Number | Reason            |
| 09/17/2012       | 1              | Original Document |
|                  |                |                   |
|                  |                |                   |
|                  |                |                   |
|                  |                |                   |
|                  |                |                   |